

Mollified Monte Carlo

hep-lat/0209053, BROWN-HET-1317

D. D. Ferrante^{* a}, J. Doll^b, G. S. Guralnik^a, D. Sabo^b.

^aPhysics Department, Brown University,
Providence, RI 02912. POBOX 1843.

^bChemistry Department, Brown University,
Providence, RI 02912.

Using a common technique [1,2] for approximating distributions [generalized functions], we are able to use standard Monte Carlo methods to compute QFT quantities in Minkowski spacetime, under phase transitions, or when dealing with coalescing stationary points.

1. Theory

We use the mollification (*approximate identity*) technique to rewrite the partition function in such a way that not only its computation, but the computation of its derivatives (Green's functions), become well defined using Monte Carlo methods, even for integrands that normally do not allow this approach.

1.1. Mollification

A mollifier, η , is a positive, $C^\infty[I]$ function, ($I \in \mathbb{R}$), with $\text{supp}(\eta) = B(0, l)$, where $l = \text{length}(I)$ and $B(0, l)$ is the ball centered at 0 with radius l and $\int_I \eta = 1$. The sequence of functions, $\eta_\epsilon = \epsilon^{-n} \eta(x/\epsilon)$ is an *approximate identity*. A *mollification* of a locally integrable function $f : U \rightarrow \mathbb{R}$, ($U \in \mathbb{R}^n$, open), is given by,

$$U_\epsilon = \{x \in U \mid \text{dist}(x, \partial U) > \epsilon\}$$

$$f_\epsilon = \eta_\epsilon * f, \quad f_\epsilon \in U_\epsilon \quad (1)$$

$$f_\epsilon(x) = \int_U \eta_\epsilon(x-y) f(y) dy \quad (2)$$

$$= \int_{B(0, \epsilon)} \eta_\epsilon(y) f(x-y) dy.$$

1.2. Properties

The properties of mollifiers, (f_ϵ), are:

1. $f_\epsilon \in C^\infty(U_\epsilon)$;
2. $f_\epsilon \rightarrow f$, almost everywhere, as $\epsilon \rightarrow 0$;

^{*}danieldf@het.brown.edu

3. If $f \in C(U)$, then $f_\epsilon \rightarrow f$ *uniformly* on compact subsets of U ; $\mathcal{E}$

4. If $1 \leq p < \infty$ and $f \in L^p_{\text{loc}}(U)$, then $f_\epsilon \rightarrow f$ in $L^p_{\text{loc}}(U)$.

It is not difficult to see that this approximation, (f_ϵ , the *mollified* version of f), can smooth oscillating functions. f_ϵ is differentiable and, not only does it converge to f when $\epsilon \rightarrow 0$ but, depending on where this convergence takes place, it can be either uniform or in L^p_{loc} .

1.3. Implementation of the Method

The object we calculate will not be the usual partition function, but its mollified version, Z_ϵ ,

$$Z_\epsilon[J] = \frac{\int \langle e^{iS(\phi) - iJ \cdot \phi} \rangle_\epsilon [d\phi]}{\int \langle e^{iS(\phi)} \rangle_\epsilon [d\phi]}, \quad (3)$$

where,

$$\langle f \rangle_\epsilon \equiv f_\epsilon \equiv \eta_\epsilon * f.$$

From equation (2) it follows that, upon choosing U appropriately, the boundary conditions² for the theory (defined here by S) are set[4].

The method is applied in two steps:

²Note that, a choice of boundary conditions [Schwinger-Dyson equation] is equivalent to choosing the measure in the path-integral.

1. Saddle-point expansion: For simplicity, the mollifier is chosen to be a Gaussian³,

$$\eta_\epsilon(\varphi) = \frac{\exp\left\{-\frac{1}{2} \varphi^\top \cdot (\epsilon^2)^{-1} \cdot \varphi\right\}}{\sqrt{(2\pi)^n \det(\epsilon^2)}}, \quad (4)$$

where ϵ^2 is the covariance matrix that defines the Gaussian distribution. Thus, a typical path-integral will look like,

$$\begin{aligned} \mathcal{J}_\epsilon &= \int \left\langle f(\varphi) e^{iS(\varphi) - iJ \cdot \varphi} \right\rangle_\epsilon [d\varphi] \\ &= f(\varphi_0) e^{iS(\varphi_0) - iJ \cdot \varphi_0} \times \\ &\times \int \frac{\exp\left\{-\frac{1}{2} \mathbf{B}^\top \cdot (\mathbb{I} + \epsilon^\top \cdot \mathbb{S}'' \cdot \epsilon)^{-1} \cdot \mathbf{B}\right\}}{\sqrt{\det(\mathbb{I} + \epsilon^\top \cdot \mathbb{S}'' \cdot \epsilon)}} [d\varphi] + \\ &+ \text{corrections}, \end{aligned}$$

where $\mathbf{B} = \epsilon \cdot \nabla S(\varphi)$ and $[\mathbb{S}'']_{ij} = \partial^2 S(\varphi) / \partial \varphi_i \partial \varphi_j$ (Hessian matrix for S).

2. Importance sampling: The Monte Carlo simulation is handled as follows: Choose an *Importance Function*, ($W(\varphi)$), and change the measure of integration accordingly,

$$\mathcal{Z}_\epsilon[J] = \frac{\int \left[\frac{\langle e^{iS(\varphi) - iJ \cdot \varphi} \rangle_\epsilon}{W(\varphi)} \right] [dW(\varphi)]}{\int \left[\frac{\langle e^{iS(\varphi)} \rangle_\epsilon}{W(\varphi)} \right] [dW(\varphi)]}. \quad (5)$$

A reasonable choice for W is given by [3] $W(\varphi) = W_\epsilon(\varphi) = |\langle e^{iS(\varphi)} \rangle_\epsilon|$. Upon a saddle-point expansion, we find,

$$W_\epsilon(\varphi) = \left| \frac{\exp\left\{iS(\varphi_0) - \frac{1}{2} \mathbf{B}^\top \cdot (\mathbb{I} + \epsilon^\top \cdot \mathbb{S}'' \cdot \epsilon)^{-1} \cdot \mathbf{B}\right\}}{\sqrt{\det(\mathbb{I} + \epsilon^\top \cdot \mathbb{S}'' \cdot \epsilon)}} \right|. \quad (6)$$

2. Applications in Physics

Two basic examples will be shown here. Both of them can be regarded as 0-dimensional [space-time] quantum field theories.

³Note that, bold symbols and bold letters, denote vector/matrix quantities.

2.1. Airy Action

The action is given by: $S(x) = ix^3/3 + itx$. Note that, $\int_{-\infty}^{\infty} \exp\{S(x)\} dx = \text{Ai}(t)$, thus, the partition function and its mollification are,


$$\mathcal{Z}[t] = \frac{\int_{-\infty}^{\infty} \exp\left\{i \frac{x^3}{3} + itx\right\} dx}{\int_{-\infty}^{\infty} \exp\left\{i \frac{x^3}{3}\right\} dx} \equiv \frac{\text{Ai}(t)}{\text{Ai}(0)}, \quad (7)$$

$$\mathcal{Z}_\epsilon[t] = \frac{\int_{-\infty}^{\infty} \left\langle \exp\left\{i \frac{x^3}{3} + itx\right\} \right\rangle_\epsilon dx}{\int_{-\infty}^{\infty} \left\langle \exp\left\{i \frac{x^3}{3}\right\} \right\rangle_\epsilon dx}. \quad (8)$$


This is a particular useful example for two reasons:

1. We can easily compare the approximated results with the exact ones,
2. Boundary Conditions: Assuming integration on the real axis, the Euclidian version of this theory does not exist, therefore working in Minkowski space is necessary.

It is easy to see that this is a highly oscillatory integrand (what makes the use of Monte Carlo techniques impracticable). However, once its mollification is calculated, the task becomes simpler (as shown below).



The importance function, ($W(x)$), is also shown for several values of the parameter ϵ :




2.2. ϕ^4 Theory


The action is, $S(\phi) = \mu \phi^2/2 + g \phi^4/4$. Thus, the mollified generating functional is given by:

$$\mathcal{Z}_\epsilon[J] = \frac{\int_{-\infty}^{\infty} \left\langle \exp \left\{ i \frac{\mu}{2} \phi^2 + i \frac{g}{4} \phi^4 - i J \phi \right\} \right\rangle_\epsilon d\phi}{\int_{-\infty}^{\infty} \left\langle \exp \left\{ i \frac{\mu}{2} \phi^2 + i \frac{g}{4} \phi^4 \right\} \right\rangle_\epsilon d\phi}.$$

As usual, its parameters, (m, g) , will determine the phase structure of the theory. The following are the plots of the 2-point Green's function in the symmetric and broken-symmetric phases, respectively.



The plots below are the importance function and the mollified integrand in the symmetric and broken symmetric phase, respectively.



It is interesting to realize that, for the broken symmetric phase, the importance function peaks exactly at the point where the mollified integrand ceases to have two concavities and begins to have only one.

3. Acknowledgements

DDF and GSG wish to acknowledge support by DOE grant DE-FG02-91ER40688 - Task D, JDD and DS wish to acknowledge support from the National Science Foundation through awards CHE-0095053 and CHE-0131114.

REFERENCES

1. L. C. Evans, *Partial Differential Equations*, AMS (1998), GSM 19.
2. M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press.
3. D. Sabo and J.D. Doll and D.L. Freeman, *Stationary Tempering and the Complex Quadrature Problem*, J. Chem. Phys., 116(9):3509-3520, March 2002.
4. S. Garcia and Z. Guralnik and G. Guralnik, *Theta Vacua and Boundary Conditions of the Schwinger-Dyson Equations*, (1996), hep-th/9612079.